




Contents lists available at ScienceDirect

Progress in Neuro-Psychopharmacology & Biological Psychiatry

journal homepage: www.elsevier.com/locate/pnpbp



Review article

Multidisciplinary approach of organic catatonia in children and adolescents may improve treatment decision making

Bertrand Lahutte ^{a,d}, Françoise Cornic ^{b,c}, Olivier Bonnot ^{a,g}, Angèle Consoli ^{a,b}, Isabelle An-Gourfinkel ^f, Zahir Amoura ^e, Frédéric Sedel ^f, David Cohen ^{a,b,g,*}

^a Département de Psychiatrie de l'Enfant et de l'Adolescent, GH Pitié-Salpêtrière, Paris, France

^b CNRS FRE 2987 Psychologie et Neurosciences Cognitives, Paris, France

^c Département de Psychiatrie, GH Sainte-Anne, Paris, France

^d Département de Psychiatrie Adulte, Service de Santé des Armées, Paris, France

^e Département de Médecine Interne, GH Pitié-Salpêtrière, Paris, France

^f Fédération de Neurologie, GH Pitié-Salpêtrière, Paris, France

^g Centre Référent Maladies Rares à Expression Psychiatrique, GH Pitié-Salpêtrière, Paris Faculté Pierre et Marie Curie, AP-HP, GHU Est, Groupe Hospitalier Pitié-Salpêtrière, 47 bd de l'Hôpital, 75013, Paris, France

ARTICLE INFO

Article history:

Received 5 January 2008

Received in revised form 27 February 2008

Accepted 27 February 2008

Available online 7 March 2008

Keywords:

Adolescence

Catatonia

Childhood

Organic conditions

Treatment decision-making

ABSTRACT

Catatonia is an infrequent but severe condition in young people. Organic diseases may be associated and need to be investigated though no specific recommendations and guidelines are available.

We extensively reviewed the literature of all the cases of organic catatonia in children and adolescents from January 1969 to June 2007. We screened socio-demographic characteristics, organic diagnosis, clinical characteristics and treatment.

We found 38 cases of children and adolescents with catatonia due to an organic condition. The catatonic syndrome occurred in 21 (57%) females and 16 (43%) males. The mean age of patients was 14.5 years (+/− 3.39) [range = 7–18 years], and three died from their condition. The organic conditions included infectious diseases ($N=10$), neurological conditions ($N=10$), toxic induced states ($N=12$) and genetic conditions including inborn errors of metabolism ($N=6$). The onset was dominantly acute, and the clinical presentation most frequently stuporous. Although benzodiazepines were recommended as primary symptomatic treatment, they were rarely prescribed. In several cases, therapeutic approach was related to organic cause (e.g., plasma exchange in lupus erythematosus; copper chelators in Wilson's disease).

Based on this review and on our own experience of catatonia in youth, we proposed a consensual and multidisciplinary diagnostic strategy to help practitioners to identify underlying organic diseases.

© 2008 Elsevier Inc. All rights reserved.

Contents

1.	Introduction	1394
2.	Method	1394
3.	Results	1394
3.1.	Characteristics of patients	1394
3.2.	Clinical characteristics	1394
3.3.	Therapeutic approaches	1395

Abbreviations: DC, David Cohen; OB, Olivier Bonnot; AC, Angèle Consoli; ZA, Zahir Amoura; FS, Frédéric Sedel; IA, Isabelle An; FC, Françoise Cornic; PANDAS, Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcal Infections; ECT, Electro-convulsive therapy; MRI, Magnetic Resonance Imaging; IEM, Inborn errors of metabolism; ADHD, Attention Deficit Hyperactive Disorder; OCD, Obsessive Compulsive Disorder; PDD, pervasive developmental disorder; NLP, neuroleptic drug; SCZ, schizophrenia; EEG, Electro-encephalography; Cbls, Cobalamin metabolism defects; MTHFR, methylene tetrahydrofolate reductase.

* Corresponding author. Centre Référent Maladies Rares à Expression Psychiatrique, GH Pitié-Salpêtrière, Paris Faculté Pierre et Marie Curie, AP-HP, GHU Est, Groupe Hospitalier Pitié-Salpêtrière, 47 bd de l'Hôpital, 75013, Paris, France. Tel.: +33 1 42 16 23 51; fax: 33 1 42 16 23 53.

E-mail address: david.cohen@psl.aphp.fr (D. Cohen).

4. Discussion	1395
4.1. Summary of the current review	1395
4.2. Clinical contribution of a multidisciplinary approach for recognition of organic underlying condition	1396
4.3. Summary of paraclinical investigations to screen organic conditions in isolated youth catatonia	1398
5. Conclusion	1398
Acknowledgement	1398
References	1398

1. Introduction

Catatonia is a rare and severe psychiatric syndrome. It is defined by the association of motor abnormalities (stupor, excitement, posturing, catalepsy, negativism, waxy flexibility, and stereotyped movements) and psychic symptoms (mutism, social withdrawal, mannerism, echolaly, verbigeration, schizophasia). Several varieties can be distinguished (Cohen et al., 2005; Taylor and Fink, 2003): stuporous catatonia, excited catatonia, malignant catatonia and psychomotor automatism. In adults, epidemiological studies using catatonia rating scales found that the prevalence of catatonia ranges from 7.6% to 38% among psychiatric inpatients. The syndrome is more frequent in female patients, is usually associated with mood disorders (Taylor and Fink, 2003), and can occur in organic conditions (Cottencin et al., 2007). In the field of child and adolescent psychiatry, few studies suggested a prevalence range from 0.6 to 17.7% (Thakur et al., 2003; Cohen et al., 2005). While the symptomatology and associated disorders are similar to those reported in the adult literature, findings differ with regard to the female-to-male ratio and the relative frequencies of associated disorders. Catatonia in children or adolescents is more frequent in boys (Takaoka and Takata, 2003) and schizophrenia is the most frequent associated diagnosis (Cornic et al., 2007). When encountered in child and adolescent clinic, the disease must lead to specific investigations, because its aetiology often reveals among psychiatric presentations, various organic diseases: neurological diseases, intoxications and metabolic conditions (Cohen et al., 1999). The stake for clinical practice resides in the potential display of a curative treatment of the underlying affection. It concerns the prognosis, by the perspective of the psychiatrist (to give the opportunity to treat the catatonic state with the treatment of the organic aetiology), but also by the perspective of the neurologist (by the recognition of psychiatric state ushering the neurologic symptoms and therefore highlighting the development of the organic condition). Besides, the diagnosis of the organic condition appears essential regarding the severity and the possible lethality of the underlying states in organic catatonia (Ainsworth, 1987; Dimitri et al., 2006).

The aims of the current study were (1) to list case reports of catatonia due to organic conditions in youths and to spot clinical characteristics and organic aetiology, and (2) to formulate recommendations and guidelines including which investigations and clinical manifestations may help determination of a cause and therefore treatment decision making.

2. Method

We conducted a literature search in the Medline data base for all reports associated with the following key-words: *catatonia* and/or *catatonic syndrome*, and *children* and/or *adolescent*. Corresponding references were then studied to determine whether cases corresponded effectively to both catatonia and organic condition criteria, and therefore could be included in this study. During the period that extended from January 1969 to June 2007, a total of 90 references were collected, among which we selected reports including medical conditions. We also performed a manual search of reference lists of the selected papers and of all reviews on catatonia in youths. In total, 30 papers mostly single case report or series were selected. We also

included three patients admitted and treated in our Department. Two were reported in a follow-up study presented at an international meeting (Cornic et al. 2006). This led to a total of 38 patients to be reviewed (Table 1). Data were extracted according to a screening collecting socio-demographic characteristics (sex, age), organic diagnosis, clinical characteristics of the catatonic syndrome according Taylor and Fink classification adapted for children and adolescent (Cohen et al. 2005), and treatment. Based on this literature review, a multidisciplinary group including experienced child psychiatrists (DC, OB, AC), one adult psychiatrist (FC), neurologists keen on epilepsy (IA) and neurometabolism (FS), and one internist (ZA), all involved in catatonic research formulated guidelines for investigation and recognition of potential organic causes in youth catatonia. These guidelines include all the causes found in the literature, but also other rare metabolic diseases that should be known by child and adolescent psychiatrists as they were identified as possible treatable causes of catatonia in youth.

3. Results

3.1. Characteristics of patients

The literature review collected 38 cases of children and adolescents with catatonia due to an organic condition reported from January 1969 to September 2007. The catatonic syndrome occurred in 21 (57%) females and 16 (43%) males. The mean age of patients was 14.5 years (+/-3.39) [range=7–18 years] and only six patients were younger than 11. Three patients died from their condition: the first had encephalitis (Ainsworth, 1987), the second had venous cortical thrombosis (Gangadhar et al., 1983) and the third had Fatal Familial Insomnia that is a rare autosomal dominant condition belonging to the prion disease group (Dimitri et al., 2006). All organic conditions encountered are listed in Table 1. They were classified as follow: infectious diseases (N=10), mainly typhoid and viral encephalitis; neurological conditions (N=10) with complex seizures and auto-immune conditions with cerebral tropism being the most frequent; toxic induced states (N=12) that may be either secondary effects of treatments (e.g., ciclosporin) or consequences of prohibited drugs (e.g., ecstasy); and finally, genetic conditions (N=6) including inborn errors of metabolism.

3.2. Clinical characteristics

Apart from few specific symptoms of the underlying organic condition (see details in Table 1), clinical characteristics of the catatonic syndrome were as follow: (1) onset was dominantly acute (96%); (2) using Taylor and Fink modified classification of catatonia, we distinguished 27 (73%) stuporous catatonia, 7 (19%) excited catatonia, 2 (5%) malignant catatonia and one case of psychomotor automatism with a progressive onset.

Regarding associated psychiatric diagnosis, in 19 cases, the only psychiatric diagnosis reported was catatonia. In the remaining 18 cases, 6 received a diagnosis of psychosis or brief psychotic disorder, 6 of psychotic depression, 3 of schizophrenia and 2 of delirium. The last patient had Attention Deficit Hyperactivity Disorder associated with Obsessive-Compulsive symptoms due to a PANDAS (Pediatric Auto-immune Neuropsychiatric Disorders Associated with Streptococcal Infections).

Table 1
Thirty height cases of organic catatonia in child ren and adolescents reported from 1969 to 2007

Diagnosis	N	Sex	Age	Clinical characteristics	Authors, year
<i>Infectious diseases (N=10)</i>					
Typhoïde	4	2M 2F	7-17- 18-18	Acute onset (N=4) Stuporous catatonia (N=2) Stuporous/excited catatonia (N=2) Fever, diarrhea (N=4)	Breckey and Kala (1977)
Viral encephalitis	4	1M 3F	14-13- 17-7	Acute onset (N=4)	Ainsworth, 1987
Infectious mononucleosis	1	1F	16	Acute onset Stuporous catatonia Neurologic signs	Rubin (1978)
Fever of unknown etiology	1	1M	17	Acute onset Stuporous catatonia Fever	Unni et al. (1995)
<i>Neurological conditions (N=10)</i>					
Seizures	3	3F	10-13- 17	Acute onset (N=3) Stuporous catatonia (N=3) Psychosis (N=2) Confusional state (N=1)	Kramer (1977), Shah and Kaplan (1980), Primavera et al. (1994)
Veinous cortical thrombosis	1	1F	18	Acute onset Stuporous catatonia	Gangadhar et al. (1983)
Neurolyupus	4	4F	13-15- 16-17	Acute onset (N=4) Stuporous catatonia (N=4) Psychotic depression (N=4)	Lanham et al. (1985), Perisse et al. (2003), Cohen et al. (2005)
Paraneoplastic limbic encephalitis	1	1F	11	Acute onset Malignant catatonia	Lee et al. (2006)
PANDAS	1	1M	11	Acute onset Stuporous/excited catatonia ADHD/OCD/No psychosis	Elia et al. (2005)
<i>Toxic induced states (N=12)</i>					
Corticotherapy	2	2M	17-11	Acute onset (N=2) Stuporous catatonia (N=2) Psychosis (N=2)	Sullivan and Dickerman (1979), Doherty et al. (1991)
Chlorphenamine maleate	1	1M	17	Acute onset Stuporous catatonia	Johnson and Lucey (1987)
Ciclosporin	1	1F	14	Acute onset Stuporous catatonia Psychotic depression	Unpublished
Ecstasy	3	1M 2F	17-17- 16	Acute onset (N=3) Stuporous catatonia (N=3) Hyponatremia (N=2)	Maxwell et al. (1993), Masi et al. (2002)
Phencyclidine	1	NR	NR	Non Specified	Baldrige and Bessen (1990)
Lithium	1	1F	16	Acute onset Stuporous catatonia	Desakar et al. (2007)
Other toxic	2	1M 1F	16-17	Stuporous/excited catatonia, acute onset (N=2), Drug/inhalant induced delirium (N=2)	Lee (1998), Lee et al. (2000)
Anaphylactic shock	1	1F	12	Acute onset Stuporous catatonia Neurologic signs	Pranzatelli et al. (1994)

Table 1 (continued)

Diagnosis	N	Sex	Age	Clinical characteristics	Authors, year
<i>Genetic conditions (N=6)</i>					
Prader-Willi syndrome	1	1M	17	Acute onset Stuporous catatonia Brief Psychotic Disorder Mild Mental Retardation	Dhossche and Bounam (1997)
Fatal Familial Insomnia	1	1M	18	Acute onset Stuporous/excited catatonia Psychotic depression Confusional state	Dimitri et al., 2006
Huntington disease	1	1M	17	Progressive onset Psychomotor automatism Schizophrenia	Unpublished, Cornic et al. (2006)
Tay-Sachs disease	1	1M	17	Progressive onset Stuporous catatonia Schizophrenia Neurologic signs	Rosebush et al. (1995)
Wilson disease	1	1M	12	Acute onset of catatonia Stuporous catatonia Neurologic history Hepatomegaly Kayser-Fleischer ring	Davis and Borde (1993)
Storage disease	1	1M	16	Acute onset Excited catatonia Schizophrenia	Unpublished Cornic et al. (2006)

NR = Not reported; *Acute defined as ≤15 days; chronic as ≥16 days; subtypes of catatonia were as follow: stuporous – excited – malignant – psychomotor automatism defined as automatic movements secondary to hallucinations being the most prevalent symptom (Cohen et al, 2005).

As for the organic examination, symptoms could be observed in 19 patients, such as fever (N=5), neurological symptoms (N=7), confusional states (N=4), hyponatremia (N=2) and hepatomegaly (N=1).

3.3. Therapeutic approaches

Among the 31 reports that indicated the therapeutic approach, psychotropic medications were used in 21 (68%) of them. Antipsychotics were used alone with 9 (29%) patients, and surprisingly – despite literature experience (Taylor and Fink, 2003; Lee et al., 2000) – major tranquilizers (benzodiazepines) in only 12 (39%): 8 (26%) associated with antipsychotics and 4 (13%) alone. ECT was realized for 11 (32%) patients. In several cases, treatment of the underlying condition was undertaken. Anti-seizures treatments were remarkable in 3 cases (Slooter et al, 2005; Shah and Kaplan, 1980; Kramer, 1977). Plasma exchanges associated or not with immuno-suppressors were so in catatonia due to auto-immune dysfunctions (Elia et al., 2006; Cohen et al., 2005; Périsse et al., 2003). Finally, copper chelators dramatically improved a 12-year-old boy with Wilson’s disease (Davis and Bordes, 1993). Regarding intoxication with ecstasy, management of hyponatremia during the early phase of the treatment in intensive care was crucial (Maxwell et al., 1993). In sum, a medical treatment focused on the organic condition was intended in 18 cases. Amidst plasma exchanges and anti-seizures, less specific medications were used (5 cases), such as large spectrum antibiotherapies, or on the other way, decrease or cessation of toxic or iatrogenic substance was realized (8 cases).

4. Discussion

4.1. Summary of the current review

The current review demonstrates that organic conditions can occur in children and adolescents catatonia and may lead to death. Although

catatonia is rare in children and adolescents, the proportion of organic causes is high for a psychiatric condition. First, Cornic et al. (2006) reported 6 organic conditions among a consecutive series of 35 patients, leading to a rate of 16%. Second, the largest review of literature of youth catatonia (Takaoka and Takata, 2003) listed 73 cases published during the period 1982–2002 including 17 cases due to an organic condition leading to a rate of 23%. Given that there is probably a bias in reporting organic cases, an estimation of the proportion of organic condition in catatonic syndromes in youth could be 15–20%. Furthermore, the review highlights that organic catatonia is not just a subtype of catatonia. Sex ratio and associated psychiatric diagnosis differs from children and adolescents non-organic catatonia, with more female than male and more acute psychosis and psychotic depression than schizophrenia in the organic group, whereas a reversal pattern is found in the non-organic group (Takaoka and Takata, 2003; Cornic et al., 2007). However, whether organic catatonia differs from non-organic ones in terms of physiopathology, response to symptomatic treatment of catatonia, prognosis and course, need to be explored although prognosis and course is linked to the accessibility of an efficient treatment towards the organic cause.

Benzodiazepines (e.g., lorazepam) or other sedative drugs and antiparkinsonism drugs (e.g., amantadine) may prove useful on catatonic manifestations and should be the first treatment option. In adult, these treatments have been shown to be helpful (Taylor and Fink, 2003). However, in the current review the poor reporting of benzodiazepine prescription highlights that the indication of these drugs in catatonia is not well known in child and adolescent psychiatric practice. In case of resistance, ECT is usually efficient on catatonia (Taylor and Fink, 2003). Finally, treatment of the underlying organic affection, when available, may be efficient as well but a rapid diagnosis is needed given the severity of most catatonic states.

4.2. Clinical contribution of a multidisciplinary approach for recognition of organic underlying condition

Fig. 1 states the general guidelines (Cornic et al., 2007) for aetiological diagnosis and treatment orientation in catatonia formu-

lated by a multidisciplinary group of physicians all involved in treatment and research in the field of catatonia.

- (1) To address organic conditions, both careful medical examination, including accurate neurological examination, and psychiatric examination are warranted to identify and rule out treatable medical disorders.
- (2) Regarding psychiatric investigations, some psychiatric manifestations are useful items for clinical orientation: type of onset, acute or insidious, mood symptoms, hallucinations, and delusions. Confusion and subtype of catatonia – stupor – should be considered, as they are the most common features in organic catatonia (Table 1). The psychiatric and medical history of the family and the patient should be collected as well as the medications used or the drug consumption. The use of catatonia rating scales to monitor symptoms should be recommended. Five validated rating scales are available in adults (Bush et al, 1996; Kruger et al, 2003; Northoff et al, 1999; Cuesta and Peralta, 2001; Lund et al, 1991) but can be used in adolescents, as well (Cohen et al, 2005).
- (3) Regarding somatic manifestations, some symptoms should be actively searched in the development of the catatonic symptomatology, or in the anamnesis of the patients, as they may orient diagnosis, through the identification of actual clinical manifestations (e.g., seizures, fever or encephalopathy), or through the existence or discovery of an evolutive organic condition (e.g., lupus erythematosus symptoms; Keiser–Fleisher corneal ring; dysmorphism; mental retardation). In summary, somatic survey will consider general examination and issuing, neurologic and ophthalmologic examination.
- (4) Paraclinical investigations should be leaded by clinical data. First line tests will complete or precise the clinical approach (see below).

Determination of organic condition type from somatic and psychiatric examination is not immediate, as pathognomonic symptoms are rare. Neurologic manifestations are omnipresent, and it should be distinguished whether they rely on neurological specific condition, or are




Fig. 1. Catatonia in children and adolescents: a multimodal framework for evaluation and treatment (PDD = pervasive developmental disorder; NLP = neuroleptic drug; SCZ = schizophrenia; ECT = electro-convulsive therapy) (adapted from Cornic et al., 2007).

Table 2

Inborn errors of metabolism that may present as catatonic states in children and adolescents: clinical characteristics and screening tests (adapted from Sedel et al, 2007)

Diseases	Psychiatric signs	Neurological signs	Systemic signs	Screening tests
<i>Treatable diseases</i>				
Urea cycle disorders	Attacks of confusion, bizarre behaviour, delusion triggered by high protein intake or situations of protein catabolism	Stroke like episodes (diplopia, hemiparesis), pyramidal signs, epilepsy, coma.	Nausea, vomiting, cephalalgia	Ammoniemia
MTHFR deficiency	Mild mental retardation, confusion, depression, psychosis	Coma, pyramidal syndrome (subacute degeneration of the cord), peripheral neuropathy, strokes.	thromboembolic events	Homocysteinemia
Cbls	Mild mental retardation, confusion, depression, psychosis	Pyramidal signs (subacute degeneration of the cord), peripheral neuropathy, optic atrophy,	retinitis pigmentosa, glomerular nephritis, thromboembolic events.	Homocysteinemia
Acute porphyrias	Episodes of confusion, psychosis, depression	Acute peripheral neuropathy, epilepsy	Intestinal problems (pain, constipation), dysautonomia, dark urines. cutaneous signs (coproporphyrin and porphyrin variegata)	Urinary porphobilinogen
Wilson's disease	Disorders of behaviour and personality, depression. Rare cases of psychosis.	Movement disorders, dysarthria	Corneal Kayser–Fleischer ring, chronic liver disease	Ceruleoplasmine, cupremia, cupruria
Cerebrotendinous xanthomatosis	Rare cases of psychosis	Cerebellar ataxia, spastic paraparesis, dementia, peripheral neuropathy, parkinsonism.	Juvenile cataract, xanthomas, chronic diarrhoea	Sterols HPLC
<i>Non treatable diseases</i>				
Metachromatic leukodystrophy	Psychosis like features (mimicks schizophrenia)	Cognitive troubles, spastic paraparesis, cerebellar ataxia, demyelinating polyneuropathy.	None	Arylsulfatase A
GM2 gangliosidosis	Episodes of psychosis, depression, mania	Lower motoneuron disease, cerebellar ataxia, pyramidal signs, dystonia, sensitive polyneuropathy	Dysautonomia	Hexosaminidases
Niemann Pick disease type C	Psychosis, depression, mania	Cognitive troubles, cerebellar ataxia, vertical oculomotor apraxia, movement disorders (dystonia, myoclonus)	Splenomegaly, hepatomegaly,	Filipin staining

included in genetic, infectious or auto-immune states. To lead this endeavour, we propose to categorize the situation, by clinical syndromic recognition.

- Neurological condition will be evoked in cases of symptoms of brain suffering (e.g. confusional states, pyramidal syndrome, and movement disorders). The spectrum of epileptic pathology may be investigated through anamnesis (of seizures, or confusional states) and accurate clinical examination, completed by electroencephalography and neuro-imaging (especially brain MRI). Encephalitis should be suspected through patent symptoms such as fever associated to neurological manifestations, but also in link with neoplastic pathologies and systemic diseases such as lupus erythematosus. Cerebral fluid analysis is therefore recommended in most cases together with search of possible germs as antibiotics and antivirals would be adjusted to the related infectious diseases.
- Auto-immune states are a crossway pathology, aside neurology and auto-immunity. Systemic lupus erythematosus will be evoked, in front of clinical symptoms such as: polyarthritis, photosensitivity, malar rash, alopecia, serositis, proteinuria and hematuria. Auto-immune investigations should be leaded in order to identify and treat as soon as possible the condition. Systemic pathologies such as lupus can be treated by a vast array of immunomodulating/suppressant drugs, including corticoids, cyclophosphamide, and hydroxychloroquine. Plasma exchanges are particularly interesting, according to the dramatic improvement of catatonia following this treatment in a recent series (Marra et al., 2008).
- Inborn errors of metabolism (IEM) may be revealed in children and adolescence by an apparently isolated psychiatric disorder. This specific focus on inborn errors of metabolism consists on the necessity of urgent specific treatments. In addition to the clinical context (testimonies of heritability related to familial history, presence of systemic or neurolo-

gical manifestations, fluctuations of symptoms triggered by catabolism, food intake, surgery etc.), psychiatric manifestations themselves are characteristic of certain types of IEM. Diseases presenting with acute attacks of confusion include urea cycle defects, homocysteine remethylation defects and porphyrias. Isolated psychiatric manifestations arising in adolescence or in a previously normal patient can be observed in patients with homocystinurias, Wilson's disease, and neuropilidosis. Catatonia, visual hallucinations and aggravation with treatments, are all atypical features that should point to an IEM. In addition, some patients have a history of mild mental retardation since childhood and behavioural or personality disorders with no clear psychiatric syndrome (Sedel et al., 2007). Regarding inherited metabolic diseases that may present with catatonic symptoms in children and adolescents, we summarized them in Table 2 and specified whether they are known as treatable or not. Treatments are variable, and may include alimentary restrictions, vitamins, symptomatic medications, or specific treatments, like copper chelators in Wilson's disease.

We consider that a systematic search of the treatable diseases is necessary even if most of these conditions are rare. Indeed, treatment at the "psychiatric stage" before the occurrence of neurological symptoms, can lead to higher frequency of reversal of symptoms (Sedel et al, 2007).

Table 3

Paraclinical investigations to screen organic conditions in isolated youth catatonia

General	Haemoglobin, blood cell count, blood chemistry (electrolytes, glucose, creatinine, blood urea, calcium, phosphate, magnesium level, liver function tests), erythrocyte sedimentation rate.
Neurological	Encephalic MRI, EEG, cerebrospinal fluid analysis (if fever)
Immunologic	Antinuclear antibodies
Toxic	Urinary drugs screening
Metabolic	Ammoniemia, homocysteinemia.

4.3. Summary of paraclinical investigations to screen organic conditions in isolated youth catatonia

As a consequence of these organic implications, the clinical encounter with a catatonic syndrome should lead to clinical and paraclinical investigations. Accurate organic diagnosis and treatment indication relies upon internists and neurologists. Unless examination is positive and suggest specific paraclinical investigations, first line ones should be sufficient to determine whether the hypothetical medical condition consists in an acute or a chronic situation, and so adapt treatment decision making. Taking into account the contribution developed above, our recommendations to screen organic conditions associated with catatonia are presented in Table 3.

5. Conclusion

This review stresses upon the fact that catatonic syndromes can be observed in children and adolescents in association with organic diseases. These are rare but severe and potentially lead to lethal conditions. Set aside the interest of orienting the aetiological diagnosis, this fact implies necessary inquiries for the clinician, in order not to neglect the perspective of treatment of the organic causal disease. The stakes are important, relying upon psychiatric symptoms reduction, but also hindering the course of metabolic or neurological diseases. Several basic investigations should be realised, lead by anamnesis, neurological and systemic symptoms, and may assist the clinician enlightened by these considerations.

Acknowledgement

Authors thank the *Fondation Wyeth pour la Santé de l'Enfant et de l'Adolescent* and the *CARPIJ* for their support in the field of severe psychiatric disorders in children and adolescents. Authors have no disclosure to declare.

References

Abczyńska M, Termińska K. Psychopathological symptoms in atypical viral hemorrhagic tick-borne encephalitis. *Psychiatr Pol* 1995;29:547–51.

Ainsworth P. A case of lethal catatonia in a 14 year-old girl. *Br J Psychiatry* 1987;150:110–2.

Baldrige EB, Bessen HA. Phencyclidine. *Emerg Med Clin North Am* 1990;8:541–50.

Breakey WR, Kala AK. Typhoid catatonia responsive to ECT. *Br Med J* 1977;2(6083):357–9.

Bush G, Fink M, Petrides G, Francis A, Catatonia I. Rating scale and standardized examination. *Acta Psychiatr Scand* 1996;93:129–36.

Cohen D, Flament M, Dubos PF, Basquin M. The catatonic syndrome in young people. *J Am Acad Child Adolesc Psych* 1999;38:1040–106.

Cohen D, Nicolas JD, Flament M, Perisse D, Dubos PF, Bonnot O, et al. Clinical relevance of chronic catatonic schizophrenia in children and adolescents: evidence from a prospective naturalistic study. *Schizophr Res* 2005;76:301–8.

Cornic F, Olliac B, Soussan N, Nicolas JD, Bonnot O, Tordjman S, et al. Clinical relevance of chronic catatonia in childhood onset schizophrenia: evidence from a 4 year F/U study. IACAPAP. (September 2006) Melbourne, Australia; 2006.

Cornic F, Consoli A, Cohen D. Catatonia in children and adolescents. *Psychiatric Annals* 2007;37:19–26.

Cottencin O, Warembourg F, de Chouly de Lenclave MB, Lucas B, Vaiva G, Goudemand M, et al. Catatonia and consultation-liaison psychiatry study of 12 cases. *Prog Neuro-Psychopharmacol Biol Psychiatry* 2007;31:1170–6.

Cuesta MJ, Peralta V. Integrating psychopathological dimensions in functional psychoses: a hierarchical approach. *Schizophr Res* 2001;52:215–29.

Davis EJ, Borde M. Wilson's disease and catatonia. *Br J Psychiatry* 1993;162:256–9.

Desarkar P, Das A, Das B, Sinha VK. Lithium toxicity presenting as catatonia in an adolescent girl. *J Clin Psychopharmacol* 2007;410–2.

Dhossche DM, Bouman NH. Catatonia in an adolescent with Prader–Willi Syndrome. *Ann Clin Psychiatry* 1997;9:247–53.

Dimitri D, Jehel L, Durr A, Levy-Soussan M, Andreux V, Laplanche JL, et al. Fatal familial insomnia presenting as psychosis in an 18-year-old man. *Neurology* 2006;67:363–4.

Doherty M, Garstin I, McClelland RJ, Rowlands BJ, Collins BJ. A steroid stupor in a surgical ward. *Br J Psychiatry* 1991;158:125–7.

Elia J, Dell ML, Friedman DF, Zimmerman RA, Balamuth N, Ahmed AA, et al. PANDAS with catatonia: a case report. Therapeutic response to lorazepam and plasmapheresis. *J Am Acad Child Adolesc Psychiatry* 2005;44:1145–50.

Gangadhar BN, Keshavan MS, Goswami U, Rao TV. Cortical venous thrombosis presenting as catatonia: a clinicopathologic report. *Clin Psychiatry* 1983;44:109–10.

Johnson J, Lucey PA. Encephalitis lethargica, a contemporary cause of catatonic stupor. A report of two cases. *Br J Psychiatry* 1987;151:550–2.

Kramer MS. Menstrual epileptoid psychosis in an adolescent girl. *Am J Dis Child* 1977;131:316–7.

Kruger S, Bagby RM, Hoffer J, Braunig P. Factor analysis of the catatonia rating scale and catatonic symptom distribution across four diagnostic groups. *Compr Psychiatry* 2003;44:472–82.

Lanham JG, Brown MM, Hughes GR. Cerebral systemic lupus erythematosus presenting with catatonia. *Postgrad Med J* 1985;61:329–30.

Lee JW. Serum iron in catatonia and neuroleptic malignant syndrome. *Biol Psychiatry* 1998;44:499–507.

Lee JW, Schwartz DL, Hallmayer J. Catatonia in a psychiatric intensive care facility: incidence and response to benzodiazepines. *Ann Clin Psychiatry* 2000;12:89–96.

Lee A, Glick DB, Dinwiddie SH. Electroconvulsive therapy in a pediatric patient with malignant catatonia and paraneoplastic limbic encephalitis. *J ECT* 2006;22:267–70.

Lund CE, Mortimer AM, Rogers D, McKenna PJ. Motor, volitional and behavioural disorders in schizophrenia. 1: assessment using the modified Rogers scale. *Br J Psychiatry* 1991;158:323–6.

Marra D, Amoura Z, Soussan N, Haroche J, Consoli A, Ghillami-Dalbin P, et al. Plasma exchange in patients with catatonia and systemic lupus erythematosus. *Psychother Psychosom* 2008;77:195–6.

Masi G, Mucci M, Floriani C. Acute catatonia after a single dose of ecstasy. *J Am Acad Child Adolesc Psychiatry* 2002;41:892.

Maxwell DL, Polkey MI, Henry JA. Hyponatraemia and catatonic stupor after taking “ecstasy”. *BMJ* 1993;307(6916):1399.

Northoff G, Koch A, Wenke J, Eckert J, Böker H, Pflug B, et al. Catatonia as a psychomotor syndrome: a rating scale and extrapyramidal motor symptoms. *Mov Disord* 1999;14:404–16.

Perisse D, Amoura Z, Cohen D, Saintigny P, Mekhloufi F, Mazet P, et al. Effectiveness of plasma exchange in an adolescent with systemic lupus erythematosus and catatonia. *J Am Acad Child Adolesc Psychiatry* 2003;42:497–9.

Pranzatelli MR, Mott SH, Pavlakis SG, Conry JA, Tate ED. Clinical spectrum of secondary parkinsonism in childhood: a reversible disorder. *Pediatr Neurol* 1994;10:131–40.

Primavera A, Fonti A, Novello P, Roccatagliata G, Cocito L. Epileptic seizures in patients with acute catatonic syndrome. *J Neurol Neurosurg Psychiatry* 1994;57:1419–22.

Rosebush PI, MacQueen GM, Clarke JT, Callahan JW, Strasberg PM, Mazurek MF. Late-onset Tay–Sachs disease presenting as catatonic schizophrenia: diagnostic and treatment issues. *J Clin Psychiatry* 1995;56:347–53.

Rubin RL. Adolescent infectious mononucleosis with psychosis. *J Clin Psychiatry* 1978;39:773–5.

Sedel F, Baumann N, Turpin JC, Saudubray JM, Cohen D. Psychiatric manifestations of inborn errors of metabolism in adolescents and adults: a clinical approach. *J Inher Metab Dis* 2007;30:631–41.

Shah A, Kaplan SL. Catatonic symptoms in a child with epilepsy. *Am J Psychiatry* 1980;137:738–9.

Slooter AJ, Braun KP, Balk FJ, van Nieuwenhuizen O, van der Hoeven J. Electroconvulsive therapy for malignant catatonia in childhood. *Pediatr Neurol* 2005;32:190–2.

Sullivan BJ, Dickerman JD. Steroid-associated catatonia: report of a case. *Pediatrics* 1979;63:677–9.

Takaoka K, Takata T. Catatonia in childhood and adolescence. *Psychiatry Clin Neurosci* 2003;57:129–37.

Taylor MA, Fink M. Catatonia in psychiatric classification: a home of its own. *Am J Psychiatry* 2003;160:1233–41.

Thakur A, Jagadsen K, Dutta S, Sinha VK. Incidence of catatonia in children and adolescents in a pediatric psychiatric clinic. *Aust N Z J Psychiatry* 2003;37:200–3.

Unni KE, Shivakumar V, Dutta TK, Chandrasekaran R. Fever of unknown cause presenting as Catatonia. *J Assoc Phys India* 1995;43:134–5.